

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

ATTY.'S DOCKET: OKAMURA=2B

In re Application of:)	Art Unit: 1646
)	
H. OKAMURA et al.)	Examiner: D. Jiang
)	
Appln. No.: 09/050,249)	Washington, D.C.
)	
Date Filed: March 30, 1998)	Confirmation No. 6601
)	
For: IFN-GAMMA PRODUCTION)	
INDUCING PROTEIN..)	

DECLARATION UNDER 37 CFR §1.132

Honorable Commissioner for Patents
U.S. Patent and Trademark Office
Customer Service Window
Randolph Building, Mail Stop
401 Dulany Street
Alexandria, VA 22314

Sir:

I, Haruki OKAMURA, declare and state as follows:

1. I am a citizen of Japan residing at Nakahozumi, 2-12-32, Ibaraki-shi, Osaka, Japan.
2. I graduated from Osaka University, Faculty of Science (Biology) and received a Doctorate of Science from Osaka University in 1976.
3. I have been working for Hyogo College of Medicine since 1976, where I am the professor of the Laboratory of Host Defenses, Institute for Advanced Medical Science.
4. I have been actively engaged in research in the field of microorganisms and cytokines, particularly mouse and human

interleukin-18 (IL-18). A copy of my curriculum vitae is attached hereto.

5. I am one of the inventors named in the above-identified patent application (hereinafter designated as "instant application") and I am thoroughly familiar with the original specification of the instant application and the presently claimed invention.

6. I understand that the examiner takes the position that the present claims are obvious/unpatentable over Nakamura et al., "Purification of a Factor which Provides a Costimulatory Signal for Gamma Interferon Production", *Infection and Immunity*, 61(1):64-70 (1993), a publication in which I am a co-author and which is hereinafter referred to as Nakamura et al. I am also thoroughly familiar with the disclosures of this publication and I hereby declare that Nakamura et al. cannot make obvious the presently claimed invention for the reasons below:

7. Summary of Nakamura et al. and relationship between Nakamura et al. and Okamura et al.

As reported in Nakamura et al., I and my co-authors found a protein factor in mouse sera, which induces a high level of interferon- γ (IFN- γ) production in the presence of interleukin-2 (IL-2), monoclonal anti-CD3 antibody (anti-CD3 MAb), or concanavalin A (ConA), and which has the following properties:

- i) The molecular weight of the protein factor was much smaller by SDS-PAGE (50 to 55 kDa) than by the molecular sieve technique (70 to 75 kDa); and
- ii) When electrophoresed on SDS-PAGE to give an apparently single protein band with a molecular size of 50 to 55 kDa, the factor lost its IFN- γ -inducing activity.

When the apparent molecular weight of the active form of a conventionally known natural killer stimulatory factor, i.e., 75 kDa NKSF/interleukin-12 (IL-12), its IFN- γ inducibility, and the synergy of IL-12, anti-CD3 MAb, or a mitogenic lectin were all found to be similar to those of the factor in Nakamura et al., I and my co-authors speculated that the factor may possibly be the natural killer stimulatory factor NKSF/IL-12. However, we could not provide proof because we were unable to obtain the factor in an isolated and purified form and in sufficient yield necessary for identifying it as either a novel substance or the natural killer stimulatory factor NKSF/IL-12.

To overcome this problem, we tried to isolate cells that produce this factor in order to facilitate the isolation and purification of the factor in a sufficient yield to perform further analysis. We screened cells, such as from established macrophage cell lines, as potential candidate cells that sufficiently express the factor. Nevertheless, despite our best efforts, we were

unsuccessful in obtaining such cells, and our research in this regard made not the slightest bit of progress.

Following the studies reported in Nakamura et al., I and my co-authors of Okamura et al., *Infection and Immunity*, "A Novel Costimulatory Factor for Gamma Interferon Induction Found in the Livers of Mice Causes Endotoxic Shock", 63(10):3966-3972 (1995), hereinafter referred to as Okamura et al., which was published after the filing date of the instant application, continued to study on how to establish the identity of the factor in Nakamura et al. and perform further analysis. For this purpose, we extensively screened cells which induce the factor from Nakamura et al.

As a result of these later studies, we unexpectedly discovered that mouse liver cells produce a substance that induces IFN- γ production and named it IGIF (Interferon-Gamma Inducing Factor). To discern the relationship between the factor from Nakamura et al. and the IGIF from Okamura et al., we re-prepared Nakamura et al.'s factor for comparison with IGIF. The re-prepared factor exhibited an IFN- γ -inducing activity in a protein species with a molecular weight of 19 kDa in addition to the one with a molecular weight of 75 to 80 kDa by molecular sieving in the presence of dithiothreitol (DTT). The activity of inducing INF- γ production, however, was observed only in the molecular species of 75 to 80 kDa in the absence of DTT. Moreover, the protein species

with a molecular weight of 75 to 80 kDa was reduced to 19 kDa on 0.1% SDS-polyacrylamide gels in the presence of DTT. In addition, the NH₂-terminal amino acid sequence of the re-prepared factor was determined to be the same as that of IGIF purified from mouse liver extracts. Based on these results, I and my co-authors of Okamura et al. concluded that *"The serum factor whose apparent molecular mass was previously found to be 75 kDa by gel filtration was shown to contain the same 18- to 19-kDa IGIF."*

8. Difficulty of obtaining the monoclonal antibody of the instant application based on Nakamura et al.

As stated above, at the time Nakamura et al. was published, I and my co-authors of Nakamura et al. had neither isolated the factor in an isolated and purified form nor identified the factor as a novel substance (even if the factor contained the IGIF later identified in Okamura et al.), and we even speculated that the factor may be the NKSF/interleukin-12 (IL-12) that induces IFN- γ production.

I hereby declare that, under these circumstances, no researcher would have considered preparing any monoclonal antibody to the factor because, first of all, I and my co-authors had not established a method in Nakamura et al. for providing sufficient amounts of the factor necessary for preparing such a monoclonal antibody. Second of all, even if monoclonal antibodies were to be obtained using the factor, nobody could have selected the desired monoclonal antibody to the factor from among the various

antibodies to the factor or possible NKSF/interleukin-12 (IL-12) present in the same protein band or fraction.

I further declare that I and my co-authors of Nakamura et al. did indeed need sufficient amounts of the factor from a source other than mouse sera in order to discern the identity of the factor; however, we were unable to establish such technique and to obtain sufficient amounts of the factor at the time.

9. Breakthrough

I and my co-inventors in the instant application consistently and diligently continued studying the factor of Nakamura et al. and, prior to the publication of Okamura et al., we found that mouse liver cells produce a mouse interleukin-18 (mIL-18) corresponding to the IGIF in Okamura et al. (hereinafter designated as "mIL-18/IGIF") and serve as a cell source for mIL-18/IGIF and we established a technique for producing sufficient amounts of mIL-18/IGIF necessary for preparing the monoclonal antibody of the instant application using recombinant DNA technology with mRNA isolated from mouse liver cells as mIL-18/IGIF-producing cells.

I declare that the above finding was the breakthrough or the key to success for obtaining the monoclonal antibody of the instant application and this is evidenced by the disclosure of Okamura et al., at page 3972, left column, lines 12 to 15:

The complementary DNA for the IGIF has recently been cloned (unpublished data). This will enable a

*sufficient supply of recombinant IGIF or **the antibody against it** for examination of its biological actions.*
(emphasis added)

I further declare that, without the above finding, one of ordinary skill in the art could never have succeeded in arriving at the monoclonal antibody of the instant application based on Nakamura et al. because the prior art and the knowledge of those of ordinary skill in the art lacked the above-mentioned key to success. Consequently, Nakamura cannot lead one of ordinary skill in the art to the presently claimed invention.

10. I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Sep. 16 2009
Date: September 16, 2009

Haruki Okamura
Haruki OKAMURA

In re of Appln. No. 09/050,249

Attachment:

Exhibit A: Curriculum vitae of Haruki OKAMURA

CURRICULUM VITAE

Dr. Haruki OKAMURA
Nakahozumi, 2-12-32, Ibaraki-shi,
Osaka, Japan

PERSONAL: Japanese Citizen
Married

EDUCATION:

1970 Graduated from Osaka University, Faculty of Science (Biology)
1976 Degree of Ph. D, Osaka University
1976 Scholarship Student, Institute for Microbial Diseases, Department of Measles,
Osaka University

BRIEF CHRONOLOGY OF EMPLOYMENT:

1976 Assistant Professor, Department of Bacteriology, Hyogo College of Medicine
1997-1999 Associate Professor, Laboratory of Host Defenses, Institute for Advanced
Medical Science, Hyogo College of Medicine
1999- Professor, Laboratory Host Defenses, Institute for Advanced Medical Science,
Hyogo College of Medicine

SOCIETIES:

Society of Immunology Japan
Society of Interferon and Cytokine Japan, Manager
Society of Inflammation Japan

PUBLICATION LIST
(Dr. Haruki OKAMURA)

- (1) Okamura, H., Tsutsui, H., Komatsu, T., Yutsudo, M., Hakura, A., Tanimoto, T., Torigoe, K., Okura, T., Nukada, Y., Hattori, K., Akita, K., Namba, M., Tanabe, F., Konishi, K., Fukuda, S., and Kurimoto, M., Cloning of a new cytokine that induces IFN- γ production by T cells. *Nature* 378, 88-91, 1995
- (2) Ushio, S., Namba, M., Okura, T., Hattori, K., Nukata, Y., Akita, K., Tanabe, F., Konishi, K., Micallef, M., Fijii, M., Torigoe, K., Tanimoto, T., Fukuda, S., Ikeda, M., Okamura, H., and Kurimoto, M., Cloning of the cDNA for Human IFN- γ -Inducing Factor, Expression in *Escherichia coli*, and Studies on the Biological Activities of the Protein lipid A. *J. Immunol.*, 156, 4274-4, 1996
- (3) Tsutsui, H., Nakanishi, K., Matsui, K., Higashino, K., Okamura, H., Miyazawa, Y., and Kaneda, K., IFN- γ Inducing Factor Up-Regulates Fas Ligand-Mediated Cytotoxic Activity of Murine Natural Killer Cell Clones. *J. Immunol.*, 157, 3367-3973, 1996
- (4) Tao, D., Ohashi, K., Tayano, T., Kurimoto, M., and Okamura, H., Interferon- γ -Inducing Factor, a Novel Cytokine, Enhances Fas Ligand-Mediated Cytotoxicity of Murine T Helper 1 Cells. *Cell. Immunol.*, 173, 230- 235, 1996
- (5) Ahn, H., Maruo, S., Tomura, M., Mu, J., Hamaoka, T., Nakanishi, K., Clark, S., Kurimoto, M., Okamura, H., and Fujiwara, H., A mechanism underlying synergy between IL-12 and IFN- γ -inducing factor (IL-18) in enhanced production of IFN- γ . *J. Immunol.*, 159, 2125-2131, 1997
- (6) Zhang, T., Kawakami, K., Qureshi, M.H., Okamura, H., Kurimoto, M., and Satoh, A., Interleukin-12(IL-12) and IL-18 synergistically induce the fungicidal activity of murine peritoneal exudate cells against *Cryptococcus neoformans* through production of gamma interferon by natural killer cells. *Infect. Immun.*, 65, 3594-3599, 1997
- (7) Stoll, S., Muller, G., Kurimoto, M., Saloga, J., Tanimoto, T., Yamauchi, H., Okamura, H., Knop, J., and Enk, A.H., Production of IL-18 (IFN- γ -inducing factor) mRNA and functional protein by murine keratinocytes. *J. Immunol.*, 159, 298-302, 1997

- (8) Gu, Y., Kuida, K., Tsutsui, H., Ku, G., Fleming, M. A., Hayashi, N., Higashino, K., Okamura, H., Nakanishi, K., Kurimoto, M., Tanimoto, T., Flavell, R. A., Sato, V., Harding, M. W., Livingston DJ., and Su MS, Activation of Interferon- γ inducing factor mediated by Interferon-1 converting enzyme. *Science*, 275, 206-209, 1997
- (9) Yoshimoto, T., Okamura, H., Tagawa, Y., Iwakura, Y., and Nakanishi, K., Interleukin-18 (IL-18) together with IL-12 inhibits IgE production by induction of IFN- γ production from activated B cells. *Proc. Natl. Acad. Sci. USA.*, 94, 3948-3953, 1997
- (10) Tsutsui, H., Matsui, K., Kawada, N., Hyodo, Y., Hayashi, N., Okamura, H., Higashino, K., and Nakanishi, K., IL-18 accounts for both TNF- α -and Fas Ligand-mediated hepatotoxic pathways in endotoxin-induced liver injury in mice. *J. Immunol.*, 159, 3961-3967, 1997
- (11) Udagawa, N., Horwood, NJ., Elliott, J., Mackay, A., Owen, J., Okamura, H., Kurimoto, M., Chambers, TJ., Martin, TJ., and Gillespie, Mt., Interleukin-18 (interferon- γ -inducing factor) is produced by osteoclasts and acts granulocyte/macrophage colony-stimulating factor and not via interferon- γ to inhibit osteoclast formation. *J. Exp. Med.*, 185, 1005-1012, 1997
- (12) Yoshimoto, T., Takeda, K., Tanaka, T., Ohkusu, K., Kashiwamura, S-I., Okamura, H., Akira, S., and Nakanishi, K., IL-12 up-regulates IL-18R expression on T cells and B cells : synergism with IL-18 for IFN- γ production. *J. Immunol.*, 161, 3400-3407, 1998
- (13) Takeda, K., Tsutsui, T., Yoshimoto, T., Adachi, O., Yoshida, N., Kishimoto, T., Okamura, H., Nakanishi, K., and Akira, S., Defective NK cell activity and Th1 response in IL-18-deficient mice. *Immunity*, 8, 383-390, 1998
- (14) Osaki, T., Peron, J. M., Cai, Q., Okamura, H., Robbins, P. D., Kurimoto, M., Lotze, M. T., and Tahara, H., IFN- γ -inducing factor/IL-18 administration mediate IFN- γ and IL-12-independent antitumor effects. *J. Immunol.*, 160, 1742-1749, 1998
- (15) Okamura, H., Tsutsui, T., Kashiwamura, S-I., Yoshimoto, T., and Nakanishi, K., Interleukin-18 (IL 18) : a novel cytokine that augments both innate and augments both innate and acquired immunity. *Adv. Immunol.*, 70, 281-312, 1998

- 6) Okamura, H., Kashiwamura, S-I., Tsutsui, H., Yoshimoto, T., and Nakanishi, K., Regulation of interferon-gamma (IFN- γ) production by IL-12 and IL-18. *Curr. Opin. Immunol.*, 10, 259-264, 1998
- 7) Osaki, T., Hashimoto, W., Gambotto, A., Okamura, H., Robbins, P. D., Kurimoto, M., Lotz, M. T., Tahara, H., Potent anti-tumor effects mediated by local expression of the mature form of interferon-gamma inducing factor, interleukin-18 (IL-18). *Gene Therapy*, 6, 808-815, 1999.
- 8) Hyodo, Y., Matsui, K., Hayashi, N., Tsutsui, H., Kashiwamura, S., Yamauchi, H., Hiroishi, K., Takeda, K., Tagawa, Y., Iwakura, Y., Kayagaki, N., Kurimoto, M., Okamura, H., Hada, T., Yagita, H., Akira, S., Nakanishi, K., Higashino, K., IL-18 up-regulates perforine-mediated NK activity without increasing perforine messenger RNA expression by binding to constitutively expressed IL-18 receptor. *J. Immunol.*, 162, 1662-1668, 1999
- 9) Rothe, H., Hausmann, A., Casteels, K., Okamura, H., Kurimoto, M., Burkart, V., Mathieu, C., and Kolb, H., IL-18 inhibits diabetes development in nonobese diabetic mice by counterregulation of Th1-dependent destructive insulinitis. *J. Immunol.*, 163, 1230-1236, 1999
- 10) Hashimoto, W., Osaki, T., Okamura, H., Robbins, P.D., Kurimoto, M., Nagata, S., Lotze, M.T., and Tahara, H. Differential antitumor effects of administration of recombinant IL-18 or recombinant IL-12 are mediated primarily by Fas-Fas ligand- and perforin-induced tumor apoptosis, respectively. *J. Immunol.*, 163, 583-9, 1999
- 1) Yogi, Y., Endoh, M., Tanaka, T., Akira, S., Okamura, H., and Nomaguchi, H. Bacteria killing by macrophages via NF-IL6 gene dependent mechanism: the susceptibility to *Mycobacterium leprae* in NF-IL6 knockout mice. *Nihon Hansenbyo Gakkai Zasshi.*, 68, 97-108, 1999
- 2) Yamamoto, T., Moriwaki, Y., Matsui, K., Takahashi, S., Tsutsui, H., Yoshimoto, T., Okamura, H., Nakanishi, K., Kurosawa, Y., Yamaguchi, S., and Sasaki, Y., Higashino K. High IL-18 (interferon-gamma inducing factor) concentration in a purine nucleoside phosphorylase deficient patient. *Arch Dis Child.*, 81, 179-80, 1999

- (23) Tsutsui, H., Kayagaki, N., Kuida, K., Nakano, H., Hayashi, N., Takeda, K., Matsui, K., Kashiwamura, S., Hada, T., Akira, S., Yagita, H., Okamura, H., and Nakanishi, K., Caspase-1-independent, Fas/Fas ligand-mediated IL-18 secretion from macrophages causes acute liver injury in mice. *Immunity*, 11, 359-67, 1999
- (24) Yoshimoto, T., Tsutsui, H., Tominaga, K., Hoshino, K., Okamura, H., Akira, S., Paul, W.E., and Nakanishi, K., IL-18, although antiallergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proc.Natl.Acad.Sci.USA.*, 96, 13962-6, 1999
- (25) Nakata, A., Tsujimura, T., Sugihara, A., Okamura, H., Iwasaki, T., Shinkai, K., Iwata, N., Kakishita, E., Akedo, H., and Terada, N., Inhibition by interleukin 18 of osteolytic bone metastasis by human breast cancer cells. *Anticancer, Res.*, 19(5B), 4131-8, 1999
- (26) Kodama, T., Matsuyama, T., Kuribayashi, K., Nishioka, Y., Sugita, M., Akira, S., Nakanishi, K., and Okamura, H., IL-18 deficiency selectively enhances allergen-induced eosinophilia in mice. *J Allergy.Clin.Immunol.*, 105(1 Pt 1), 45-53, 2000
- (27) Tominaga, K., Yoshimoto, T., Torigoe, K., Kurimoto, M., Matsui, K., Hada, T., Okamura, H., and Nakanishi, K., IL-12 synergizes with IL-18 or IL-1beta for IFN-gamma production from human T cells. *Int. Immunol.*, 12, 151-60, 2000
- (28) Ohkusu, K., Yoshimoto, T., Takeda, K., Ogura, T., Kashiwamura, S., Iwakura, Y., Akira, S., Okamura, H., and Nakanishi, K., Potentiality of interleukin-18 as a useful reagent for treatment and prevention of leishmania major infection. *Infect. Immun.*, 68, 2449-56, 2000
- (29) Ida, A., Tsuji, Y., Muranaka, J., Kanazawa, R., Nakata, Y., Adachi, S., Okamura, H., and Koyama, K., IL-18 in pregnancy; the elevation of IL-18 in maternal peripheral blood during labour and complicated pregnancies. *J. Reprod. Immunol.*, 47, 65-74, 2000
- (30) Tsutsui, H., Matsui, K., Okamura, H., and Nakanishi, K., Pathophysiological roles of interleukin-18 in inflammatory liver diseases. *Immunol, Rev.*, 174, 192-209, 2000

- (31) Chang, J.T., Segal, B.M., Nakanishi, K., Okamura, H. and Shevach, E.M., The costimulatory effect of IL-18 on the induction of antigen-specific IFN- γ production by resting T cells is IL-12 dependent and is mediated by up-regulation of the IL-12 receptor 2 subunit. *Eur.J. Immunol.*, 30, 1113-1119, 2000
- (32) Fujimori, Y., Takatsuka, H., Takemoto, Y., Hara, H., Okamura, H., Nakanishi, K., Kakishita, E. Elevated interleukin (IL)-18 levels during acute graft-versus-host disease after allogeneic bone marrow transplantation. *British J. Haematology.*, 109, 652-657, 2000
- (33) Yamanaka, K., Tanaka, M., Tsutsui, H., Kupper, T.S., Asahi, K., Okamura, H., Nakanishi, K., Suzuki, M., Kayagaki, N., Black, R.A., Miller, D.K., Nakashima, K., Shimizu, M. and Mizutani, H. Skin-specific caspase-1 transgenic mice show cutaneous apoptosis and pre-shock condition with a high serum level of IL-18. *J. Immunol.*, 165, 997-1003, 2000
- (34) Yoshimoto, T., Mizutani, H., Tsutsui, H., Noben-Trauth, N., Yamanaka, K-I., Tanaka, M., Izumi, S., Okamura, H., Paul, W.E. and Nakanishi, K. IL-18 induction of IgE: dependence on CD4+T cells, IL-4 and STAT6. *Nature Immunol.*, 1, 132-137, 2000
- (35) Chikano, S., Sawada, K., Shimoyama, T., Kashiwamura, S-I., Sugihara, A., Sekikawa, K., Terada, N., Nakanishi, K. and Okamura, H. IL-18 and IL-12 induce intestinal inflammation and fatty liver in mice in an IFN- γ dependent manner. *Gut*, 47, 779-786, 2000
- (36) Shigehara, K., Shijubo, N., Ohmichi, M., Yamada, G., Takahashi, R., Okamura, H., Kurimoto, M., Hiraga, Y., Tatsuno, T., Abe, S. and Sato, N. Increased levels of interleukin-18 in patients with pulmonary sarcoidosis. *Am. J. Respir. Crit. Care. Med.*, 162, 1979-1982, 2000
- (37) Tanaka, F., Hashimoto, W., Okamura, H., Robbins, P.D., Lotze, M.T. and Tahara, H. Rapid generation of potent and tumor-specific cytotoxic T lymphocytes by interleukin 18 using dendritic cells and natural killer cells. *Cancer Res.*, 60, 4838-4844, 2000
- (38) Yamada, G., Shijubo, N., Shigehara, K., Okamura, H., Kurimoto, M. and Abe, S. Increased levels of circulating interleukin-18 in patients with advanced tuberculosis. *Am. J. Respir. Crit. Care. Med.*, 161, 1786-1789, 2000

- (39) Kuhara, T., Iigo, M., Itoh, T., Ushida, Y., Sekine, K., Terada, N., Okamura, H., and Tsuda, H. Orally administered lactoferrin exerts an antimetastatic effect and enhances production of IL-18 in the intestinal epithelium. *Nutr. Cancer*, 38, 192-199, 2000
- (40) Eberl, M., Beck, E., Coulson, P.S., Okamura, H., Wilson, R.A., and Mountford, A.P. IL-18 potentiates the adjuvant properties of IL-12 in the induction of a strong Th1 type immune response against a recombinant antigen. *Vaccine*, 18, 2002-2008, 2000
- (41) Makiishi-Shimobayashi, C., Tsujimura, T., Iwasaki, T., Yamada, N., Sugihara, A., Okamura, H., Hayashi, S., and Terada, N. Interleukin-18 up-regulates osteoprotegerin expression in stromal/osteoblastic cells. *Biochem. Biophys. Res. Commun.*, 281, 361-366
- (42) Seki, E., Tsutsui, H., Nakano, H., Tsuji, N., Hoshino, K., Adachi, O., Adachi, K., Futatsugi, S., Kuida, K., Takeuchi, O., Okamura, H., Fujimoto, J., Akira, S., and Nakanishi, K. Lipopolysaccharide-induced IL-18 secretion from murine Kupffer cells independently of myeloid differentiation factor 88 that is critically involved in induction of production of IL-12 and IL-1 β . *J. Immunol.*, 166, 2651-2657, 2001
- (43) Shigehara, K., Shijubo, N., Ohmichi, M., Takahashi, R., Kon, S., Okamura, H., Kurimoto, M., Hiraga, Y., Tatsuno, T., Abe, S., and Sato, N. IL-12 and IL-18 are increased and stimulate IFN- γ production in sarcoid lungs. *J. Immunol.*, 166, 642-649, 2001
- (44) Sugawara, S., Uehara, A., Nochi, T., Yamaguchi, T., Ueda, H., Sugiyama, A., Hanzawa, K., Kumagai, K., Okamura, H., and Takada, H. Neutrophil proteinase 3-mediated induction of bioactive IL-18 secretion by human oral epithelial cells. *J. Immunol.*, 167, 6568-6575, 2001
- (45) Tsuji, Y., Tamaoki, T.H., Hasegawa, A., Kashiwamura, S., Iemoto, A., Ueda, H., Muranaka, J., Adachi, S., Furuyama, J., Okamura, H., and Koyama, K. Expression of interleukin-18 and its receptor in mouse ovary. *Am. J. Reprod. Immunol.*, 46, 349-357, 2001
- (46) Adachi, K., Tsutsui, H., Kashiwamura, S., Seki, E., Nakano, H., Takeuchi, O., Takeda, K., Okumura, K., Van, Kaer, L., Okamura, H., Akira, S., and Nakanishi, K. Plasmodium berghei infection in mice induces liver injury by an IL-12- and toll-like receptor/myeloid differentiation factor 88-dependent mechanism. *J. Immunol.*, 167, 5928-5934, 2001

- (47) Nomaguchi, H., Jahan, N., Mandal, B.C., Yogi, Y., Kawatsu, K., Yoshizawa, Y., Okamura, H., and Makino, M. IL-12 and IL-18 synergistically induce the bactericidal activity of murine peritoneal cells against *M. leprae*. *Nihon, Hansenbyo, Gakkai, Zasshi*, 70, 113-119, 2001
- (48) Ogura, T., Ueda, H., Hosohara, K., Tsuji, R., Nagata, Y., Kashiwamura, S., and Okamura, H. Interleukin-18 stimulates hematopoietic cytokine and growth factor formation and augments circulating granulocytes in mice. *Blood*, 98, 2101-2107, 2001
- (49) Itoi, H., Fujimori, Y., Tsutsui, H., Matsui, K., Futatsugi, S., Okamura, H., Hara, H., Hada, T., Kakishita, E., and Nakanishi, K. Fas ligand-induced caspase-1-dependent accumulation of interleukin-18 in mice with acute graft-versus-host disease. *Blood*, 98, 235-237, 2001
- (50) Shida, K., Shiratori, I., Matsumoto, M., Fukumori, Y., Matsuhisa, A., Kikkawa, S., Tsuji, S., Okamura, H., Toyoshima, K., and Seya, T. An alternative form of IL-18 in human blood plasma: complex formation with IgM defined by monoclonal antibodies. *J. Immunol.*, 166, 6671-6679, 2001
- (51) Nakanishi, K., Yoshimoto, T., Tsutsui, H., and Okamura, H. Interleukin-18 is a unique cytokine that stimulates both Th1 and Th2 responses depending on its cytokine milieu. *Cytokine, Growth, Factor, Rev.*, 12, 53-72, 2001
- (52) Wang, W., Tanaka, T., Okamura, H., Sugita, M., Higa, S., Kishimoto, T., and Suemura, M. Interleukin-18 enhances the production of interleukin-8 by eosinophils. *Eur. J. Immunol.*, 1010-1016, 2001
- (53) Nakanishi, K., Yoshimoto, T., Tsutsui, H., and Okamura, H. Interleukin-18 regulates both Th1 and Th2 responses. *Annu. Rev. Immunol.*, 423-474, 2001
- (54) Makiishi-Shimobayashi, C., Tsujimura, T., Iwasaki, T., Yamada, N., Sugihara, A., Okamura, H., Hayashi, S., and Terada, N. Interleukin-18 up-regulates osteoprotegerin expression in stromal/osteoblastic cells. *Biochem. Biophys. Res. Commun.*, 281, 361-366, 2001

- (55) Seki, E., Tsutsui, H., Nakano, H., Tsuji, N., Hoshino, K., Adachi, O., Adachi, K., Futatsugi, S., Kuida, K., Takeuchi, O., Okamura, H., Fujimoto, J., Akira, S., and Nakanishi, K. Lipopolysaccharide-induced IL-18 secretion from murine Kupffer cells independently of myeloid differentiation factor 88 that is critically involved in induction of production of IL-12 and IL-1 β . *J. Immunol.*, 166, 2651-2657, 2001
- (56) Shigehara, K., Shijubo, N., Ohmichi, M., Takahashi, R., Kon, S., Okamura, H., Kurimoto, M., Hiraga, Y., Tatsuno, T., Abe, S., and Sato, N. IL-12 and IL-18 are increased and stimulate IFN- γ production in sarcoid lungs. *J. Immunol.*, 166, 642-649, 2001
- (57) Sugawara, S., Uehara, A., Nochi, T., Yamaguchi, T., Ueda, H., Sugiyama, A., Hanzawa, K., Kumagai, K., Okamura, H. and Takada, H. Neutrophil proteinase 3-mediated induction of bioactive IL-18 secretion by human oral epithelial cells. *J. Immunol.*, 167, 6568-6575, 2001.
- (58) Tsuji, Y., Tamaoki, T.H., Hasegawa, A., Kashiwamura, S., Iemoto, A. Ueda, H., Muranaka, J., Adachi, S., Furuyama, J., Okamura, H. and Koyama, K. Expression of interleukin-18 and its receptor in mouse ovary. *Am. J. Reprod. Immunol.*, 46, 349-357, 2001.
- (59) Adachi, K., Tsutsui, H., Kashiwamura, S., Seki, E., Nakano, H., Takeuchi, O., Takeda, K., Okumura, K., Van, Kaer, L., Okamura, H., Akira, S. and Nakanishi, K. Plasmodium berghei infection in mice induces liver injury by an IL-12- and toll-like receptor/myeloid differentiation factor 88-dependent mechanism. *J. Immunol.*, 167, 5928-5934, 2001.
- (60) Nomaguchi, H., Jahan, N., Mandal, B.C., Yogi, Y., Kawatsu, K., Yoshizawa, Y., Okamura, H. and Makino, M. IL-12 and IL-18 synergistically induce the bactericidal activity of murine peritoneal cells against *M. leprae*. *Nihon Hansenbyo Gakkai Zasshi*, 70, 113-119, 2001.
- (61) Ogura, T., Ueda, H., Hosohara, K., Tsuji, R., Nagata, Y., Kashiwamura, S. and Okamura H. Interleukin-18 stimulates hematopoietic cytokine and growth factor formation and augments circulating granulocytes in mice. *Blood*, 98, 2101-2107, 2001.
- (62) Mori, I., Hossain, M.J., Takeda, K., Okamura, H., Imai, Y., Kohsaka, S., and Kimura, Y. Impaired microglial activation in the brain of IL-18-gene-disrupted mice after neurovirulent influenza A virus infection. *Virology*, 287, 163-170, 2001.

- (63) Itoi, H., Fujimori, Y., Tsutsui, H., Matsui, K., Futatsugi, S., Okamura, H., Hara, H., Hada, T., Kakishita, E. and Nakanishi, K. Fas ligand-induced caspase-1-dependent accumulation of interleukin-18 in mice with acute graft-versus-host disease. *Blood*, 98, 235-237, 2001.
- (64) Shida, K., Shiratori, I., Matsumoto, M., Fukumori, Y., Matsuhisa, A., Kikkawa, S., Tsuji, S., Okamura, H., Toyoshima, K. and Seya, T. An alternative form of IL-18 in human blood plasma: complex formation with IgM defined by monoclonal antibodies. *J. Immunol.*, 166, 6671-6679, 2001.
- (65) Wang, W., Tanaka, T., Okamura, H., Sugita, M., Higa, S., Kishimoto, T. and Suemura, M. Interleukin-18 enhances the production of interleukin-8 by eosinophils. *Eur. J. Immunol.*, 31, 1010-1016, 2001.
- (66) Makiishi-Shimobayashi, C., Tsujimura, T., Iwasaki, T., Yamada, N., Sugihara, A., Okamura, H., Hayashi, S. and Terada, N. Interleukin-18 up-regulates osteoprotegerin expression in stromal/osteoblastic cells. *Biochem. Biophys. Res. Commun.*, 281, 361-366, 2001.
- (67) Seki, E., Tsutsui, H., Nakano, H., Tsuji, N., Hoshino, K., Adachi, O., Adachi, K., Futatsugi, S., Kuida, K., Takeuchi, O., Okamura, H., Fujimoto, J., Akira, S. and Nakanishi, K. Lipopolysaccharide-induced IL-18 secretion from murine Kupffer cells independently of myeloid differentiation factor 88 that is critically involved in induction of production of IL-12 and IL-1beta. *J. Immunol.*, 166, 2651-2657, 2001.
- (68) Shigehara, K., Shijubo, N., Ohmichi, M., Takahashi, R., Kon, S., Okamura, H., Kurimoto, M., Hiraga, Y., Tatsuno, T., Abe, S. and Sato, N. IL-12 and IL-18 are increased and stimulate IFN-gamma production in sarcoid lungs. *J. Immunol.*, 166, 642-649, 2001.
- (69) Minagawa, K., Tsuji, Y., Ueda, H., Koyama, K., Tanizawa, K., Okamura, H. and Hashimoto-Tamaoki, T. Possible correlation between high levels of IL-18 in the cord blood of pre-term infants and neonatal development of periventricular leukomalacia and cerebral palsy. *Cytokine*, 17, 164-170, 2002.

- (70) Singh, R.P., Kashiwamura, S., Rao, P., Okamura, H., Mukherjee, A. and Chauhan, V.S. The role of IL-18 in blood-stage immunity against murine malaria *Plasmodium yoelii* 265 and *Plasmodium berghei* ANKA. *J. Immunol.*, 168, 4674-4681, 2002.
- (71) Kokai, M., Kashiwamura, S., Okamura, H., Ohara, K. and Morita, Y. Plasma interleukin-18 levels in patients with psychiatric disorders. *J. Immunother.*, 25(Suppl.1), S68-71, 2002.
- (72) Futani, H., Okayama, A., Matsui, K., Kashiwamura, S., Sasaki, T., Hada, T., Nakanishi, K., Tateishi, H., Maruo, S. and Okamura, H. Relation between interleukin-18 and PGE2 in synovial fluid of osteoarthritis: a potential therapeutic target of cartilage degradation. *J. Immunother.*, 25, S61-64, 2002.
- (73) Iwasaki, T., Yamashita, K., Tsujimura, T., Kashiwamura, S., Tsutsui, H., Kaisho, T., Sugihara, A., Yamada, N., Mukai, M., Yoneda, T., Okamura, H., Akedo, H. and Terada, N. Interleukin-18 inhibits osteolytic bone metastasis by human lung cancer cells possibly through suppression of osteoclastic bone-resorption in nude mice. *J. Immunother.*, 25(Suppl.1), S52-60, 2002.
- (74) Kimura-Shimmyo, A., Kashiwamura, S., Ueda, H., Ikeda, T., Kanno, S., Akira, S., Nakanishi, K., Mimura, O. and Okamura, H. Cytokine-induced injury of the lacrimal and salivary glands. *J. Immunother.*, 25(Suppl.1), S42-51, 2002.
- (75) Kashiwamura, S., Ueda, H. and Okamura, H. Roles of interleukin-18 in tissue destruction and compensatory reactions. *J. Immunother.*, 25(Suppl.1), S4-S11, 2002.
- (76) Lotze, M.T., Tahara, H. and Okamura, H. Interleukin-18 as a novel, distinct, and distant member of the interleukin-1 family promoting development of the adaptive immune response: the interleukin-18 issue of the Journal of Immunotherapy. *J. Immunother.*, 25(Suppl.1), S1-S3, 2002.
- (77) Kuribayashi, K., Kodama, T., Okamura, H., Sugita, M., and Matsuyama, T. Effects of post-inhalation treatment with interleukin-12 on airway hyper-reactivity, eosinophilia and interleukin-18 receptor expression in a mouse model of asthma. *Clin. Exp. Allergy*, 32, 641-649, 2002.

- (78) Uchimura, T., Motomiya, Y., Okamura, H., Hashiguchi, T., Miura, M., Uji, Y., Iwamoto, H. and Maruyama, I. Marked increases in macrophage colony-stimulating factor and interleukin-18 in maintenance hemodialysis patients: comparative study of advanced glycation end products, carboxymethyllysine and pentosidine. *Nephron*, 90, 401-407, 2002.
- (79) Yamada, N., Niwa, S., Tsujimura, T., Iwasaki, T., Sugihara, A., Futani, H., Hayashi, S., Okamura, H., Akedo, H. and Terada, N. Interleukin-18 and interleukin-12 synergistically inhibit osteoclastic bone-resorbing activity. *Bone*, 30, 901-908, 2002.
- (80) Hosohara, K., Ueda, H., Kashiwamura, S., Yano, T., Ogura, T. Marukawa, S. and Okamura, H. Interleukin-18 induces acute biphasic reduction in the levels of circulating leukocytes in mice. *Clin. Diagn. Lab. Immunol.*, 9, 777-783, 2002.
- (81) Fujimori, Y., Yoshimoto, T., Matsui, K., Tsutsui, H., Okamoto, T., Kashiwamura, S., Hada, T., Okamura, H., Kakishita, E., Hara, H. and Nakanishi, K. Increased expression of interleukin-18 receptor on T lymphocytes in patients with acute graft-versus-host disease after allogeneic bone marrow transplantation. *J. Interferon Cytokine Res.*, 22, 751-754, 2002.
- (82) Naito, Y., Tsujino, T., Fujioka, Y., Ohyanagi, M., Okamura, H. and Iwasaki, T. Increased circulating interleukin-18 in patients with congestive heart failure. *Heart*, 88, 296-297, 2002.
- (83) Tokmadzic, V.S., Tsuji, Y., Bogovic, T., Laskarin, G., Cupurdija, K., Strbo, N., Koyama, K., Okamura, H., Podack, E.R. and Rukavina, D. IL-18 is present at the maternal-fetal interface and enhances cytotoxic activity of decidual lymphocytes. *Am. J. Reprod. Immunol.*, 48, 191-200, 2002.
- (84) Yamashita K, Iwasaki T, Tsujimura T, Sugihara A, Yamada N, Ueda H, Okamura H, Futani H, Maruo S, and Terada N. Interleukin-18 inhibits lodging and subsequent growth of human multiple myeloma cells in the bone marrow. *Oncol. Rep.*, 9, 1237-1244.
- (85) Hayashi, H. Inoue, Y, Tsutsui H, Okamura H, Nakanishi K, Onozaki K. (2003) TGF-beta down-regulates IFN-gamma production in IL-18 treated NK cell line LNKSE6. *Biochem. Biophys. Res. Commun.*, 300, 980-985, 2002.

- (86) Hashimoto, W., Tanaka, F., Robbins, P.D., Taniguchi, M. and Okamura, H., Lotze, M.T. and Tahara, H. Natural killer, but not natural killer T, cells play a necessary role in the promotion of an innate antitumor response induced by IL-18. *Int. J. Cancer*, 103, 508-513, 2003.
- (87) Kaneda, M., Kashiwamura, S., Ueda, H., Sawada, K., Sugihara, A., Terada, N., Kimura-Shimmyo, A., Fukuda, Y., Shimoyama, T. and Okamura, H. Inflammatory Liver Steatosis Caused by IL-12 and IL-18. *J. Interferon Cytokine Res.*, 23, 155-162, 2003.
- (88) Moriwaki, Y., Yamamoto, T., Shibutani, Y., Aoki, E., Tsutsumi, Z., Takahashi, S., Okamura, H., Koga, M., Fukuchi, M. and Hada, T. Elevated levels of interleukin-18 and tumor necrosis factor-alpha in serum of patients with type 2 diabetes mellitus: relationship with diabetic nephropathy. *Metabolism*, 52, 605-608, 2003.
- (89) Kawasaki, D., Tsujino, T., Morimoto, S., Fujioka, Y., Naito, Y., Okumura, T., Masutani, M., Shimizu, H., Yuba, M., Ueda, A., Ohyanagi, M., Kashiwamura, S., Okamura, H. and Iwasaki, T. Usefulness of circulating interleukin-18 concentration in acute myocardial infarction as a risk factor for late restenosis after emergency coronary angioplasty. *Am. J. Cardiol.*, 91, 1258-1261, 2003.
- (90) Roles of caspase-1 in *Listeria* infection in mice. Tsuji, N.M., Tsutsui, H., Seki, E., Kuida, K., Okamura, H., Nakanishi, K. and Flavell, R.A. *Int. Immunol.*, 16, 335-343, 2004.
- (91) Okamoto, T., Yamada, N., Tsujimura, T., Sugihara, A., Nishizawa, Y., Ueda, H., Kashiwamura, S., Tsutsui, H., Futani, H., Maruo, S., Okamura, H. and Terada, N. Inhibition by interleukin-18 of the growth of Dunn osteosarcoma cells. *J. Interferon Cytokine Res.*, 24, 161-167, 2004.
- (92) Oku, H., Tsuji, Y., Kashiwamura, S.I., Adachi, S., Kubota, A., Okamura, H. and Koyama, K. Role of IL-18 in pathogenesis of endometriosis. *Hum. Reprod.*, 19, 709-714, 2004.
- (93) Itoi, H., Fujimori, Y., Tsutsui, H., Matsui, K., Hada, T., Kakishita, E., Okamura, H., Hara, H. and Nakanishi, K. Differential upregulation of interleukin-18 receptor alpha chain between CD4+ and CD8+ T cells during acute graft-versus-host disease in mice. *J. Interferon Cytokine Res.*, 24, 291-296, 2004.

- (94) Kanno, T., Nagata, T., Yamamoto, S., Okamura, H. and Nishizaki, T. Interleukin1-18 stimulates synaptically released glutamate and enhances postsynaptic AMPA receptor responses in the CA1 region of mouse hippocampal slices. *Brain Res.*, 25, 190-193, 2004.
- (95) Itoi, H., Fujimori, Y., Tsutsui, H., Matsui, K., Sugihara, A., Terada, N., Hada, T., Kakishita, E., Okamura, H., Hara, H. and Nakanishi K. Involvement of interleukin-18 in acute graft-versus-host disease in mice. *Transplantation*, 15, 1245-1250, 2004.
- (96) Sekiyama, A (Reprint)., Ueda, H., Kashiwamura, S., Takeda, M., and Okamura, H. iNOS is involved in regulation of IL-18 levels in plasma. *Nitric Oxide-Biology and Chemistry*, (AUG 2004) Vol. 11, No. 1, 69-69
- (97) Yoshikawa, R (Reprint)., Yanagi, H., Hashimoto-Tamaoki, T., Morinaga, T., Nakano, Y., Noda, M., Fujiwara, Y., Okamura, H., and Yamamura, T. Gene expression in response to anti-tumour intervention by polysaccharide-K (PSK) in colorectal carcinoma cells. *ONCOLOGY REPORTS*, (DEC 2004) Vol. 12, No. 6, 1287-1293.
- (98) Nakanishi, K (Reprint)., Itoi H., Fujimori, Y., Tsutsui, H., Matsui, K., Sugihara, M., Terada, N., Hada T., Kakishita E., Okamura, H. and Hara, H. Involvement of interleukin-18 in acute graft-versus-host disease in mice. *TRANSPLANTATION*, (15 NOV 2004) Vol. 78, No. 9, 1245-1250.
- (99) Sawada, K (Reprint)., Kashiwamura, S., Okamura, H., Ohnishi, K., Fukunaga, K., Saniabadi, A. and Hada T. Enhanced efficacy of interferon-alpha and ribavirin combination by selective depletion of infected granulocytes and monocytes: Studies in patients with high plasma HCV viraemia. *GASTROENTEROLOGY*, (APR 2004) Vol. 126, No. 4, Supp. [2], A694-A694.
- (100) Nishizaki, T (Reprint)., Kanno, T., Nagata, T., Yamamoto, S., and Okamura, H. Interleukin-18 stimulates synaptically released glutamate and enhances postsynaptic AMPA receptor responses in the CA1 region of mouse hippocampal slices. *BRAIN RESEARCH*, (25 JUN 2004) Vol. 1012, No. 1-2, 190-193.
- (101) Nakanishi, K (Reprint)., Itoi, H., Fujimori, Y., Tsutsui, H., Matsui, K., Hada, T., Kakishita, ., Okamura, H. and Hara, H. Differential upregulation of interleukin-18

receptor alpha chain between CD4+ and CD8+ T cells during acute graft-versus-host disease in mice. JOURNAL OF INTERFERON AND CYTOKINE RESEARCH, (MAY 2004) Vol. 24, No. 5, 291-296.

- (102) Terada, N (Reprint), Okamoto, T., Yamada, N., Tsujimura, T., Sugihara, A., Nishizawa, Y., Ueda, H., Kashiwamura, S I., Tsutsui, H., Futani, H., Maruo, S., and Okamura, H. Inhibition by interleukin-18 of the growth of Dunn osteosarcoma cells. JOURNAL OF INTERFERON AND CYTOKINE RESEARCH, (MAR 2004) Vol. 24, No. 3, 161-167.
- (103) Tsuji, Y (Reprint), Oku H., Kashiwamura, S I., Adachi, S., Kubota, A., Okamura, H., and Koyama, K. Role of IL-18 in pathogenesis of endometriosis HUMAN REPRODUCTION, (MAR 2004) Vol. 19, No. 3, 709-714.
- (104) Flavell R A (Reprint), Tsuji, N M., Tsutsui, H., Seki, E., Kuida, K., Okamura, H. and Nakanishi, K. Roles of caspase-1 in Listeria infection in mice INTERNATIONAL IMMUNOLOGY, (FEB 2004) Vol. 16, No. 2, 335-343.
- (105) Ueda, H., and Okamura, H. Inhibition of inflammation by IL-18 Rinsho Men'eki (2004), 41(4), 455-459
- (106) Ogura, T., and Ohtsubo, Y. Antiviral effects of aciclovir injectable formulation (AcirovecR and ZoviraxR for i.v. infusion) against herpes simplex virus type 1 in vitro and in vivo. Japanese Pharmacology and Therapeutics, (2004) Vol. 32, No. 6, 349-352.
- (107) Kotani, J (Reprint), Ueno, N., Kashiwamura, S., Ueda, H., Okamura, H., Tsuji, N M., Hosohara, K., and Marukawa, S. Role of interleukin 18 in nitric oxide production and pancreatic damage during acute pancreatitis. SHOCK, (DEC 2005) Vol. 24, No. 6, 564-570.
- (108) Yoshida, K (Reprint), Okamura, H., Amorim B R., Ozaki, A., Tanaka, H., Morimoto, H., and Haneji, T. Double-stranded RNA-dependent protein kinase is required for bone calcification in MC3T3-E1 cells in vitro. EXPERIMENTAL CELL RESEARCH, (15 NOV 2005) Vol. 311, No. 1, 117-125.
- (109) Tsujino, T (Reprint), Kawasaki, D., Morimoto, S., Masai, M., Masutani, M., Ohyanagi, M., Kashiwamura, S., Okamura, H., and Masuyama T, Plasma interleukin-18

concentration: a novel marker of myocardial ischemia rather than necrosis in humans. CORONARY ARTERY DISEASE, (NOV 2005) Vol. 16, No. 7, 437-441.

- (110) Yamada, N (Reprint)., Tsujimura, T., Ueda, H., Hayashi, S I., Ohyama, H., Okamura, H., and Terada N. Down-regulation of osteoprotegerin production in bone marrow macrophages by macrophage colony-stimulating factor. CYTOKINE, (21 AUG 2005) Vol. 31, No. 4, 288-297.
- (111) Morimoto, H (Reprint)., Ozaki, A., Okamura, H., Yoshida, K., Kitamura, S., and Haneji, T. Okadaic acid induces tyrosine phosphorylation of I kappa B alpha that mediated by PKR pathway in human osteoblastic MG63 cells. MOLECULAR AND CELLULAR BIOCHEMISTRY, (AUG 2005) Vol. 276, No.1-2, 211-217.
- (112) Sawada, K (Reprint)., Kashiwamura, S., Okamura, H., Ohnishi, K., Fukunaga, K., Hirata, I., and Saniabadi, A. Selective granulocyte and monocyte apheresis as a new adjunct to enhance the efficacy of interferon-alpha plus ribavirin in patients with high plasma hepatitis C virus. DIGESTIVE AND LIVER DISEASE, (JUL 2005) Vol. 37, No. 7, 515-521.
- (113) Okamura, H (Reprint)., Nakatani-Okuda, A., Ueda, H., Kashiwamura, S., Sekiyama, A., Kubota, A., Fujita, Y., Adachi, S., Tsuji, Y., and Tanizawa, T. Protection against bleomycin-induced lung injury by IL-18 in mice. AMERICAN JOURNAL OF PHYSIOLOGY-LUNG CELLULAR AND MOLECULAR PHYSIOLOGY, (AUG 2005) Vol. 289, No. 2, L280-L287.
- (114) Sekiyama, A (Reprint)., Ueda, H., Kashiwamura, S., Sekiyama, R., Takeda, M., Rokutan, K., and Okamura, H. A stress-induced, superoxide-mediated caspase-1 activation pathway causes plasma IL-18 upregulation. IMMUNITY, (JUN 2005) Vol. 22, No. 6, 669-677.
- (115) Fujimori, Y (Reprint)., Yamaguchi, N., Fujibayashi, Y., Kasumoto, I., Okamura, H., Nakanishi, K., and Hara, H. Interferon-gamma production by human cord blood monocyte-derived dendritic cells. ANNALS OF HEMATOLOGY, (JUL 2005) Vol. 84, No. 7, 423-428.
- (116) Sekiyama, A (Reprint)., Ueda, H., Kashiwamura, S., and Okamura, H. Stress induced,

superoxide mediated caspase-1 activation causes plasma cytokines' elevation.

BIOLOGICAL PSYCHIATRY, (15 APR 2005) Vol. 57, No. 8, Supp. [S], 92S-92S. MA 336.

- (117) Ukai, K (Reprint)., Nakagawa, T., Ohyama, T., Koida, M., and Okamura, H. Effects of the synthesized growth hormone releasing peptide, KP-102, on growth hormone release in sodium glutamate monohydrate-treated low growth rats (Vol 59, p. 705, 1996). LIFE SCIENCES, (22 APR 2005) Vol. 76, No. 23, . 2751-+.
- (118) Emoto, N (Reprint)., Rohman, M S., Nonaka, H., Okura, R., Nishimura, M., Yagita, K., van der Horst G T J., Matsuo, M., Okamura, H., and Yokoyama, M. Circadian clock genes directly regulate expression of the Na⁺/H⁺ exchanger NHE3 in the kidney. KIDNEY INTERNATIONAL, (APR 2005) Vol. 67, No. 4, 1410-1419.
- (119) Sekiyama, A (correspondence)., Nishida, K., Kawai, K., Ueda, H., Kashiwamura, S., Okamura, H., Teshima-Kondo, S., and Rokutan, K. IL-18., a cytokine translates a stress into medical science. Journal of Medical Investigation, (Nov 2005) Vol. 52, No. SUPPL., 236-239.
- (120) Nakatani, O., Akemi, T., Takakuni-Ueda, H., Kashiwamura, S., Sekiyama, A., Okamura, H (correspondence)., Kubota, A., Fujita, Y., Adachi, S., and Tsuji, Y. Protection against bleomycin-induced lung injury by IL-18 in mice. American Journal of Physiology - Lung Cellular and Molecular Physiology, (Aug 2005) Vol. 289, No. 2 33-2, L280-L287.
- (121) Okamura, H., IL-1 family (IL-1alpha/beta, IL-1Ra, IL-18), IL-16, IL-17. Nippon rinsho. Japanese journal of clinical medicine, (2005 Apr) Vol. 63 Suppl 4, 226-33.
- (122) Okamura, H., Kashiwamura, S., and Ueda, H. Interleukin-18 in pathogenesis of rheumatoid arthritis. Nippon rinsho. Japanese journal of clinical medicine, (2005 Jan) Vol. 63 Sulpp. 1, 175-80.
- (123) Nakatani-Okuda, A., Ueda, H., Kashiwamura, S., Sekiyama, A., Kubota, A., Fujita, Y., Adachi, S., Tsuji, Y., Tanizawa, T., and Okamura, H. Protection against bleomycin-induced lung injury by IL-18 in Mice. American journal of physiology. Lung cellular and molecular physiology 2005), 33(2), L280-L287, 39 refs.

- (124) Fujioka, Y (Reprint)., Okumura, T., Morimoto, S., Masai, M., Sakoda, T., Tsujino, T., Kashiwamura, S., Okamura, H., and Ohyanagi, M. Chylomicron remnants stimulate release of interleukin-1 beta by THP-1 cells. JOURNAL OF ATHEROSCLEROSIS AND THROMBOSIS, (FEB 2006) Vol. 13, No. 1, 38-45.
- (125) Yoshikawa, R (Reprint)., Yanagi, H., Shen C S., Fujiwara, Y., Noda, M., Yagyu, T., Gega, M., Oshima, T., Yamamura, T., Okamura, H., Nakano, Y., Morinaga, T., and Hashimoto-Tamaoki, T. ECA39 is a novel distant metastasis-related biomarker in colorectal cancer. WORLD JOURNAL OF GASTROENTEROLOGY, (28 SEP 2006) Vol. 12, No. 36, 5884-5889.
- (126) Okamura, H., Light activates adrenal gland via suprachiasmatic nucleus NEUROSCIENCE RESEARCH, (2006) Vol. 55, Supp. [1], S44-S44.
- (127) Hirata, J (Reprint)., Kotani, J., Kirita, M., Ueda, H., Kashiwamura, S., Okamura, H., and Marukawa, S. The similarity and difference of the role(s) in human neutrophil apoptosis between IL-18 and lipopolysaccharide (LPS). SHOCK (JUN 2006) Vol. 25, Supp. [1], 67-67.
- (128) Kuhara, T (Reprint)., Yamauchi, K., Tamura, Y., and Okamura, H. Oral administration of lactoferrin increases NK cell activity in mice via increased production of IL-18 and type IIFN in the small intestine. JOURNAL OF INTERFERON AND CYTOKINE RESEARCH, (JUL 2006) Vol. 26, No. 7, 489-499.
- (129) Sawai, H (Reprint)., Hori, M., Tsuji, Y., Okamura, H., and Koyama, K. Bone morphogenetic protein-2 counterregulates interleukin-18 mRNA and protein in MC3T3-E1 mouse osteoblastic cells. CONNECTIVE TISSUE RESEARCH, (MAY-JUN 2006) Vol. 47, No. 3, 124-132.
- (130) Yamada, N (Reprint)., Nakamura, Y., Ohyama, H., Nakasho, K., Nishizawa, Y., Okamoto, T., Futani, H., Yoshiya, S., Okamura, H., and Terada N. Effect of interleukin-18 on metastasis of mouse osteosarcoma cells. CANCER IMMUNOLOGY IMMUNOTHERAPY, (SEP 2006) Vol. 55, No. 9, 1151-1158.
- (131) Okamura, H (Reprint)., Ueda, H., Kashiwamura, S I., Sekiyama, A., Ogura, T., and Gamachi, N. Production of IL-13 in spleen cells by IL-18 and IL-12 through generation

of NK-like cells. CYTOKINE, (21 FEB 2006) Vol. 33, No. 4, 179-187.

- (132) Kasuga, M (Reprint)., Inoue, H., Ogawa, W., Asakawa, A., Okamoto, Y., Nishizawa, A., Matsumoto, M., Teshigawara, K., Matsuki, Y., Watanabe, E., Hiramatsu, R., Notohara, K., Katayose, K., Okamura, H., Kahn C R., Noda, T., Takeda, K., Akira, S., and Inui, A. Role of hepatic STAT3 in brain-insulin action on hepatic glucose production. CELL METABOLISM, (APR 2006) Vol. 3, No. 4, 267-275.
- (133) Sekiyama, A (Reprint)., Ueda, H., Kashiwamura, S., Nishida, K., Yamaguchi, S., Sasaki, H., Kuwano, Y., Kawai, K., Teshima-Kondo, S., Rokutan, K., and Okamura, H A role of the adrenal gland in stress-induced up-regulation of cytokines in plasma. JOURNAL OF NEUROIMMUNOLOGY, (FEB 2006) Vol. 171, No. 1-2, 38-44.
- (134) Sekiyama, A (Reprint Author)., Nishida, K., Yamaguchi, S., Kawai, K., Yamamoto, Y., Kashiwamura, S., Ueda, H., Rokutan, K., Teshima-Kondo, S., and Okamura, H. Opposed effect of stress on inflammation is mediated by stress-induced cytokines. Biological Psychiatry, (APR 15 2006) Vol. 59, No. 8, Suppl. S, 164S.
- (135) Sekiyama, A (Reprint Author)., Nishida, K., Yamaguchi, S., Kawai, K., Yamamoto, Y., Kashiwamura, S., Ueda, H., Rokutan, K., Teshima-Kondo, S., and Ura H. Opposed effect of stress on inflammation is mediated by stress-induced cytokines. Biological Psychiatry, (APR 15 2006) Vol. 59, No. 8, Suppl. S, 78S.
- (136) Ogura, T, Kataoka, H., Ohtsubo, Y., Ueda, H., and Okamura, H. In vitro comparative experiments of aciclovir injectables, Acirovec injection and Zovirax for I.V. infusion, on anti-herpes virus activity. Igaku to Yakugaku (2006), 55(6), 893-900.
- (137) Nagai, N., Fukuhata, T., Liu Ying., Ito, Y., Kawakami, Y., Ikeda, T., Mimura, O., and Okamura, H. Relation between cell damage and expression of iNOS mRNA in human lens epithelial cell line SRA 01/04 with treatment by interferon- γ and lipopolysaccharide. Atarashii Ganka (2006), 23(3), 383-386.
- (138) Ueno, N., Hosohara, K., Kotani, J (correspondence)., and Marukawa, S. Role of interleukin 18 in nitric oxide production and pancreatic damage during acute pancreatitis. Shock, (Jan 2006) Vol. 24, No. 6, 564-570.

- (139) Ito, Y (Reprint)., Nagai, N., and Okamura, H. Involvement of interleukin 18 in cataract development in hereditary cataract UPL rats. JOURNAL OF BIOCHEMISTRY, (NOV 2007) Vol. 142, No. 5, 597-603.
- (140) Ishiguro, K (Reprint)., Oku, H., Ogawa, Y., Iwaoka, E., Kunitomo, M., Ueda, H., and Okamura, H. Involvement of inducible nitric oxide synthase in blood flow decrease in vein induced by hen-egg white lysozyme. BIOLOGICAL & PHARMACEUTICAL BULLETIN, (JUL 2007) Vol. 30, No. 7, 1324-1328.
- (141) Okamura, H (Reprint)., Li Wen., Kashiwamura, S., Ueda, H., and Sekiyama, A. Protection of CD8+ T cells from activation-induced cell death by IL-18. JOURNAL OF LEUKOCYTE BIOLOGY, (JUL 2007) Vol. 82, No. 1, 142-151.
- (142) Fujimori, Y (Reprint)., Fujibayashi, Y., Kasumoto, I., Kai, S., Hara, H., Okamura, H., Tsutsui, H., Ogawa, H., and Nakanishi, K. Interleukin-18 regulates T helper 1 or 2 immune responses of human cord blood CD4(+) V alpha 24(+)Vss11(+) natural killer T cells. EXPERIMENTAL AND MOLECULAR PATHOLOGY, (APR 2007) Vol. 82, No. 2, 220-226.
- (143) Nishida, K (Reprint)., Kamizato, M., Kiyoshi, M., Kawai, T., Teshima-Kondo, S., Okamura, H., Sekiyama, A., and Rokutan, K. IL-18 regulates both constitutive and stress-induced expression of Muc2 gene in the mouse colon. GASTROENTEROLOGY, (APR 2007) Vol. 132, No. 4, Supp. [2], A394-A394.
- (144) Sekiyama, A (Reprint)., Nishida, K., Kasahara, E., Haruyasu, U., Rokutan, K., and Okamura, H. Interleukin-18 is involved in alteration of hypothalamic-pituitary-adrenal axis activity by stress. BIOLOGICAL PSYCHIATRY, (15 APR 2007) Vol. 61, No. 8, Supp. [S], 123S-123S. MA 398.
- (145) Okamura, H (Reprint)., Seino, H., Ueda, H., Kokai, M., Tsuji, N M., Kashiwamura, S., and Morita, Y. IL-18 mediates the formation of stress-induced, histamine-dependent gastric lesions. AMERICAN JOURNAL OF PHYSIOLOGY-GASTROINTESTINAL AND LIVER PHYSIOLOGY, (JAN 2007) Vol. 292, No. 1, G262-G267.
- (146) Nagai, N., Song, Y., Fukuhata, T., Ito, Y., and Okamura, H. Involvement of interleukin-18 in cataract formation in UPL rat. Atarashii Ganka (2007), 24(6), 815-818.

- (147) Seino, H., Morita, Y., and Okamura, H. IL-18 mediates the formation of stress-induced, histamine-dependent gastric lesions. *American Journal of Physiology - Gastrointestinal and Liver Physiology*, (Jan 2007) Vol. 292, No. 1, G262-G267.
- (148) Miyoshi, K., Obata, K., Kondo, T., Noguchi, K (Reprint)., and Okamura, H. Interleukin-18-Mediated Microglia/Astrocyte Interaction in the Spinal Cord Enhances Neuropathic Pain Processing after Nerve Injury. *JOURNAL OF NEUROSCIENCE*, (26 NOV 2008) Vol. 28, No. 48, 12775-12787.
- (149) Kotani, J (Reprint)., Hirata, J., Aoyama, M., Marukawa, S., Usami, M., Kashiwamura, S., Ueda, H., Okamura, H., and Kuroda, Y. A ROLE FOR IL-18 IN HUMAN NEUTROPHIL APOPTOSIS. *SHOCK* (DEC 2008) Vol. 30, No. 6, 628-633.
- (150) Kashiwamura, S (Reprint)., Hosotani, Y., Kimura-Shimmyo, A., Sekiyama, A., Ueda, H., Ikeda, T., Mimura, O., and Okamura, H. Interleukin-18 prevents apoptosis via PI3K/Akt pathway in normal human keratinocytes. *JOURNAL OF DERMATOLOGY*, (AUG 2008) Vol. 35, No. 8, 514-524.
- (151) Terada, N (Reprint)., Watanabe, H., Hata, M., Ueda, H., Yamada, N., Yamanegi, K., Ohyama, H., Kakihana, M., Okamura, H., and Nakasho, K. Transdifferentiation into biliary ductular cells of hepatocytes transplanted into the spleen. *PATHOLOGY*, (2008) Vol. 40, No. 3, 272-276.
- (152) Kasahara Emiko (Reprint)., Sekiyama Atsuo., Okuda Akiko., Okamura Haruki. Disturbance of plasma level of stress-related cytokines in mental Disorders. *BIOLOGICAL PSYCHIATRY*, (1 APR 2008) Vol. 63, No. 7, Supp. [S], 98S-98S. MA 297.
- (153) Yamada, N (Reprint)., Nishio, S., Ohyama, H., Yamanegi, K., Nakasho, K., Hata M., Nakamura, Y., Fukazawa, S., Futani, H., Yoshiya, S., Ueda, H., Taniguchi, M., Okamura, H., and Terada, N. Enhanced suppression of pulmonary metastasis of malignant melanoma cells by combined administration of alpha-galactosylceramide and interleukin-18. *CANCER SCIENCE*, (JAN 2008) Vol. 99, No. 1, 113-120.
- (154) Yamanegi, K (Reprint)., Yamanegi, K (Reprint)., Yamane, J., Hata, M., Ohyama, H., Yamada, N., Kato-Kogoe, N., Nakasho, K., Terada, N., Futani, H., and Okamura, H.

Sodium valproate, a histone deacetylase inhibitor, decreases the secretion of soluble Fas by human osteosarcoma cells and increases their sensitivity to Fas-mediated cell death. JOURNAL OF CANCER RESEARCH AND CLINICAL ONCOLOGY, (JUL 2009), Vol. 135, No. 7, 879-889.

- (155) Sekiyama, A (Reprint)., Kasahara, E., Inoue, M., Sekiyama, R., and Okamura, H. Loss of Association between Levels of Interleukin (IL)-6 and IL-1b in Plasma in Major Depressive Disorder. BIOLOGICAL PSYCHIATRY, (15 APR 2009) Vol. 65, No. 8, Supp. [S], 233S-234S. MA 765.
- (156) Sekiyama, A (Reprint)., Kasahara, E., Inoue, M., Sekiyama, A (Reprint)., Sekiyama, R., and Okamura, H. Stress is Visible., Distinct Levels of Cytokines and Chemokines in Plasma After Psychological and Physical Loads. BIOLOGICAL PSYCHIATRY, (15 APR 2009) Vol. 65, No. 8, Supp. [S], 109S-110S. MA 366.
- (157) Suzuki, T (Reprint Author)., Acharya B., Terao, S., Yamamoto, S., Shima, H., Okamura, H., and Gotoh, A. THE ANITUMOR EFFECT OF gamma Delta T CELLS AGAINST BLADDER CANCER CELLS IS ENHANCED FOLLOWING UPREGULATION OF NKG2D LIGANDS BY VALPROIC ACID. Journal of Urology, (APR 2009) Vol. 181, No. 4, Suppl. S, 348.
- (158) Ogura, T., and Okamura, H. Comparative studies of Epirubicin hydrochloride for injection [SAWAI] and Farmorubicin for injection on in vivo and in vitro antitumor activities. Japanese Pharmacology & Therapeutics (2009), 37(3), 223-230.